

General Emergency Protocol

asthma

ACTION PLAN

Green Zone: Routine Therapy

If someone with asthma is:

- breathing easily
- not coughing or wheezing
- not short of breath
- able to work and play easily

Avoid these possible asthma triggers

Dust mites, mold, chemicals, changes in stress, temperature, tobacco smoke, cockroaches, and pets.

QUICK RELIEF Medicines:

Take 5 minutes prior to exercise or unavoidable trigger exposure.

Yellow Zone: Step Up Therapy

If someone with asthma is:

- coughing
- wheezing
- short of breath
- having difficulty breathing during activity
- feeling chest tightness

Take QUICK RELIEF Medicines

Encourage the person to:

- rest in a comfortable position, but not lying down
- relax and take slow deep breaths.

Monitor Symptoms:

- ☐ If symptoms RESOLVE within 15 minutes, the person with asthma may return to normal duties.
- ☐ If symptoms PERSIST or return within a few hours, follow red zone directions.

Red Zone: Get Help NOW!

If:

- medicine is not helping
- breathing is very difficult
- breathing difficulty limits mild activity
- speaking makes the person short of breath

Take QUICK RELIEF Medicines

- If red zone symptoms persist, call 911.
- If symptoms do not improve, repeat previous QUICK RELIEF doses and watch the person closely until ambulance arrives.

DO NOT try and treat severe symptoms yourself.